

_RTS S_NT_ MÒNIC_

ANTARCTICA. TIME OF CHANGE
MIREYA MASÓ



**ANTARCTICA
TIME OF CHANGE
MIREYA MASÓ**

BARCELONA 2010



_RTS S_NT_ MÒNIC_



Edition

ACTAR/Arts Santa Mònica
Texts©the authors
Translations©the translators
Images and Pictures©the photographers
This edition©Actar/Arts Santa Mònica- Ministry of Culture
and the Media, Government of Catalonia

Distributed by
Actar, Barcelona - New York
Part of ActarBirkhäuser
www.actar.com

ActarBirkhäuser
Distribution
Barcelona - Basel - New York
www.actarbirkhauser.com

Roca i Batlle 2
E-08023 Barcelona
T +34 93 417 49 43
F +34 93 418 67 07
office@actar-d.com

Viaduktstrasse 42
CH -4051 Basel
T +41 61 2050 777
F +41 61 2050 792
sales@birkhauser.ch

151 Grand Street, 5th floor
New York, NY 10013, USA
T +1 212 966 2207
F +1 212 966 2214
officeusa@actar-d.com

Bookshops of the Government of Catalonia
www.gencat.cat/publicacions

ISBN 978-84-393-8284-3 (Arts Santa Mònica)
ISBN 978-84-92861-26-2 (ACTAR)
DL B-29841-2010

Arts Santa Mònica
La Rambla, 7
08002 Barcelona
Tel.:+34 93 316 28 10
www.artssantamonica.cat

Actar
Barcelona/New York
info@actar.com
www.actar.com

Government of Catalonia Ministry of Culture and the Media

Minister for Culture and the Media
Joan Manuel Tresserras

Secretary General
Lluís Noguera

Secretary for Culture
Eduard Voltas

Secretary for the Media
Carles Mundó

Entitat Autònoma de Difusió Cultural
Conxita Oliver

Arts Santa Mònica

Director
Vicenç Altaió

Arts Ambit
Manuel Guerrero

Science Ambit
Josep Perelló

Communication Ambit
David Vidal

General Coordination
Marta Garcia
Fina Duran

Administration
Cristina Güell

Public Relations
Alicia Gonzalez

Editor
Cinta Massip

Press and Communication
Neus Purri
Cristina Suau

Production and Coordination
Lurdes Ibarz
Ester Martínez
Arantza Morlius

Technical Area
Xavier Roca
Eulàlia Garcia

Administrative officer
Pep Xaus
Carles Ferry

Arts Santa Mònica gratefully acknowledges the support of the Corporació Catalana de Mitjans Audiovisuals, of *La Vanguardia*, *Vilaweb* and *Mau Mau* as collaborating media, and of the Fundació Bosch i Gimpera of the Universitat de Barcelona, and the INCOM of the Universitat Autònoma de Barcelona as collaborating academic institutions.



Iaac
Institute for
advanced
architecture
of Catalonia

do you commit to action
on climate change?



The Diatoms of Antarctica and their Potential Roles in Nanotechnology

Richard Gordon, Andrzej Witkowski, Ille Christine Gebeshuber & Claire S. Allen

Biological nanotechnology and man-made nanotechnology currently still differ substantially (Abdel-aal & Ille C. Gebeshuber, 2010). Whereas in biological systems, intricate hierarchical structures are built with integrated functionalities on various length scales (from the nanometer scale to the micrometer scale to the millimeter scale), engineers still struggle to control shape or functionalities in their nanomachinery. We are just at humble beginnings of the development of hierarchical engineered structures. Diatoms, with their amazing structures and functionalities, with their ability for cell-division once a day or even more often, and with their ability to produce silica even in Antarctica, where the water temperature is below 0°C, can serve as inspiration in human technology, yielding sustainable materials, structures and processes. Diatoms have become fascinating to the industrial world over the past two decades primarily because of their ability to create solid structures of many types at a wide range of size scales. Some of them have motors with no moving parts that run at 99.9% efficiency (Gordon, 1987), and all produce oil from which we could run our cars in a sustainable way (Ramachandra et al., 2009). Antarctic diatoms accomplish these feats under conditions that are extreme, such as low temperature, at high salinity embedded in or attached to ice (Janech et al., 2006; Krell et al., 2008), and often under high ultraviolet radiation from the sun (Helbling et al., 1996; Skerratt et al., 1998; Hernando & Ferreyra, 2005; Wulff et al., 2008). They survive the Antarctic winter for 6 months with little or no light (Wulff et al., 2008) and many remain viable even when frozen into sea ice or buried in sediment, with some able to germinate after many years (Davis, 1972; Zgurovskaya, 1977; Hollibaugh, Seibert & Thomas, 1981; Ligowski, Godlewski & A. Łukowski, 1992). Antarctic diatoms thus rightfully belong to those microorganisms that are called extremophiles (Sterrenburg et al., 2007). As we shall show, these properties may prove useful in applications of diatom nanotechnology.

Diatoms are both the cause and possible solution to the present global warming crisis, though global warming has at least one positive side, as farming has contributed to it for 5000 years and may have postponed the next ice age (Ruddiman, 2005). Those of us who put up with or escape Canadian winters sometimes schizophrenically yearn for global warming. The role of diatoms is quite simple: much of the crude oil that we take out of the ground comes from diatoms, and can still be seen within the shells of diatoms long dead. If we were to switch to live diatoms, perhaps living in solar panels and genetically engineered to secrete their oil, or even gasoline (petrol) itself (Ramachandra et al., 2009), then there would be no net increase in carbon dioxide: they would take up as much as we put into the atmosphere on burning their oil. Furthermore, there would be no need to switch to new kinds of automobile engines, such as electric or hydrogen based.

Nanotechnology has come to mean all things tiny that we manufacture. One Nobel Prize winning physicist, Robert B. Laughlin, calls this infatuation the production of “*nanobaubles*, fascinating and beautiful structures that develop spontaneously at small scales but have no known use except as entertainment.... While our knowledge of the nanoscale is exploding almost incomprehensibly at the moment, nearly all of it is deeply unimportant.... The idea that nanoscale objects ought to be controllable is so compelling it blinds a person to the overwhelming evidence that they cannot be” (Laughlin, 2005).

The ordinary approach to nanotechnology is to create it by lithography (Sargent, 2005; Dupas, Houdy & Lahmani, 2007; Reisner, 2008; Gebeshuber, 2009) or chemical reaction, often biochemical (Goodsell, 2004; Rehm, 2006; Renugopalakrishnan & Lewis, 2006; Gazit, 2007; Papazoglou & Parthasarathy, 2007). Here our focus is on diatom nanotechnology, in which one uses living organisms to grow desired nanostructures rather than trying to build them ourselves:

“Diatoms... generate nanostructured silica microshells (frustules) with thousands of species-specific morphologies. Sustained reproduction of a particular diatom species

can yield enormous numbers of frustules with similar 3D morphologies. Such intricate, genetically precise, and massively parallel 3D self-assembly under ambient conditions lies well beyond the current capabilities of synthetic micro- and nanofabrication” (Weatherspoon et al., 2007).

In an offhand way, one of us started the field of diatom nanotechnology when invited to give his first talk at an engineering conference, and not knowing what to say, decided to suggest that diatoms could make things engineers wanted (Gordon & Aguda, 1988). It has since burgeoned into a large effort by many people, tying together biologists, called diatomists, with industrialists (Gordon, Sterrenburg & Sandhage, 2005; Kroth et al., 2007; Allison et al., 2008; Kröger & Poulsen, 2008; Bozarth, Maier & Zauner, 2009). The plans are wide sweeping, including “everything but the kitchen sink” in miniature form:

“Actuator, antenna, bar, bearing, cantilever, capsule, catalyst, cone, cube, cylinder, die, diffraction grating, disk, fiber, filler, filter, funnel, gear, heat exchanger, hinge, honeycomb, insulator, lens, lever, light pipe, magnet, membrane, mesh, mirror, mixer, motor, needle, nozzle, piston engine, plate, prism, pulley, pump, reactor, refraction grating, relay, rocket, rotor, sensor, separator, sieve, sphere, spiral, spring, substrate, switch, syringe, tag, tetrahedron, tetrakaidecahedron, transducer, tube, turbine engine, valve, wedge, wheel” (Gordon, 2010).

One might agree with Laughlin, because this perhaps \$100 million investment so far has yet to yield a single commercial product (Gordon, 2010). But perhaps in time it will. Certainly, diatoms figure in an art form little appreciated by the art world: the careful arrangement of diatoms to produce patterns and pictures only visible with a microscope (Nagy, 2002; Matthias Burba, 2008; Kemp, 2009). Diatoms, including those from Antarctica, have appeared on postage stamps (Edlund, 2009).

Diatom nanotechnology is intimately tied to the field of morphogenesis, in which one tries to figure out how organisms get their shapes (i.e., morphology). This is one of the basic, unsolved problems of biology. One of the refreshing aspects of all this attention to diatoms is that, in order to manipulate them, most industrialists and their scientist collaborators believe it would help if we could figure out how diatoms actually create their silica structures. Thus basic science is getting an enormous boost, and many pieces of the puzzle have been found, if not yet put together into a coherent mechanism (Gordon, 2008). So playing with all these “nanobaubles” just might lead to breakthroughs in our understanding of life in general: “Lots of money is sloshing around, and great fortunes are being made and lost.... The allure of travelling in such a wild and lawless place is the ever-present possibility of making a serendipitous discovery of great importance” (Laughlin, 2005).

The basics of diatom morphology have already been covered in this book (Scharek, 2010). Diatom assemblages of the Antarctic and the Austral Islands are unique in many respects. Firstly they inhabit either sea ice or cold to very cold waters (Medlin & Priddle, 1990). Secondly their habitats are subject to very strong winds. The latter phenomenon results in mixing of the assemblages. Hence marine and likewise terrestrial assemblages are composed of both marine and freshwater forms (Witkowski, Riaux-Gobin & Daniszewska-Kowalczyk, 2010). In addition the whole area determined as Southern Ocean is isolated from surrounding areas by a strong hydrologic barrier – the Polar Front. The existence of the Polar Front significantly reduces the exchange of the organisms with the other more northern geographic regions. Under such harsh environmental conditions peculiar diatom assemblages have developed composed of diatom species that are endemic for the Antarctic (Southern Ocean) as outlined by the Polar Front. Included in this assemblage are numerous genera, but the most interesting are *Fragilariopsis* and *Eucampia*. They are indicative either for marine ice or for conditions in the water column. Due to this they also play very important roles as indicators of climate change (Zielinski & Gersonde, 1997).

The diatom assemblages we studied originated from fairly deep stations (up to 50m). They were high in species number and predominantly composed of benthic species,

FIGURE 6




FIGURE 7

FIGURE 6. A girdle band 0.4 mm in diameter from a "giant" centric Antarctic diatom, *Arachnoidiscus* sp. It is decorated with much smaller, adhering *Cocconeis* pennate diatoms shown in the closeup. Scale bars are 0.3 mm and 0.03 mm, respectively.

FIGURE 7. The valve view of a giant diatom *Trigonium arcticum* 0.5 mm wide, and a *Coscinodiscus asteromphalus* 0.25 mm wide, found in Antarctic waters, the latter showing an array of pores, which could be used for a nanofilter.

FIGURE 8





FIGURE 9

FIGURE 8. The two valves of *Corethron* at either end of a long cylinder of girdle bands. The valve to the right has a set of long spines attached to the rim, the valve to the left has similar spines, but alternating with them is a series of finer hooked spines. Scale bar 50 μ m. Image used with permission, © R.M. Crawford and F. Hinz, from (Gebeshuber & Crawford, 2006).

FIGURE 9. A cell of *Corethron* with the old valve (V1) to the left showing the base of a number of long spines and the new valve (V2) to the right. In the new valve, the spines are still oriented parallel to the cylinder of the girdle-bands. Scale bar 20 μ m. Image used with permission, © R.M. Crawford and F. Hinz, from (Gebeshuber & Crawford, 2006).

FIGURE 10




FIGURE 10. Surface view of a valve similar to that in the right in Figure 8, showing insertion of the long spines at the edge of the valve. Scale bar 10 μm . Image used with permission. (© R.M. Crawford and F. Hinz, from Gebeshuber & Crawford, 2006). b) Closeup.

FIGURE 11




FIGURE 11. "Filtration applications of diatoms. (a) A filtration column is packed with diatom frustules. Large molecules will pass through the column relatively quickly, while smaller molecules will be able to enter the frustules via their pores and will thus be eluted at a much lower rate. (b) Biosensor filter: in a typical application (e.g. monitoring blood glucose), receptor molecules are contained (either free or fixed to the support, as shown here) within a chamber capped by a diatom frustule. Small molecules may enter via the pores in the frustule and bind to the receptors, eliciting a signal. Larger molecules capable of disrupting the signal (e.g. proteases) are prevented from entering the chamber." From (Parkinson & Gordon, 1999) with permission from Elsevier.

though planktonic and ice diatoms were also observed. This is only possible due to very high transparency of the water column allowing deep penetration of sunshine. So far little is known about ecology and species composition of Antarctic littoral/sublittoral (near shore) diatom assemblages. Research effort is mainly focused on sea ice and planktonic diatoms as they play important roles in primary production of the Antarctic waters. Hence numerous studies were focused on survival strategies of Antarctic diatoms. Interestingly some of the planktonic diatoms form wintering stages (Fryxell, 1994) while others survive in sea ice (Cunningham & Leventer, 1998).

This imbalance in attention to sea ice and planktonic diatoms on one hand and marine benthic diatoms of the Antarctic on the other is difficult to understand as the latter forms offer much more diverse and novel morphologies, which are possible to observe nowhere else in the world's ocean. This provides exciting research opportunities first of all for diatomologists dealing with biodiversity and taxonomy (new species and higher taxa) and to nanotechnologists (new ultrastructures).

Let's now have a look at some diatom parts, to see why engineers are intrigued with their potential. We'll use examples from our own work. A simple one is the circle or hoop, which comes from the silica girdle bands found around most diatoms. What is remarkable in this example is the high degree of perfection of circularity, as if it were "precisely machined" (Figure 6), but we have no lathes that small. Of course, a ring is but a component in a potential nanomachine, but that is the current state of the art: no one has yet assembled any device with moving diatom parts. A nanorobot (nanobot) consisting of a diatom shell propelled by bacteria motors has been proposed for carrying drugs inside our bloodstream, and one might imagine the optical diffracting powers of diatom shells to be put to work in arrays fancier than movable micromirrors (Link & Zimmerman, 2007).

Diatom morphologies are amazing not only because of their geometric variety and perfection but also because of their size range. Antarctic diatoms range in size from just 1 to 2 microns in length and width (e.g. *Fragilariopsis cylindrus*, Figure 5) to 3 to 4 mm in length and only a few microns wide (*Thalassiothrix* spp., *Trichotoxon* spp. and *Entopyla* spp.) and up to several hundred microns diameter (*Arachnoidiscus* spp.: Figure 6, and *Coscinodiscus* spp.: Figure 7 and Figure 5). Even within a single species the size ratio of individuals can be almost an order of magnitude (10x)!

Some of the largest diatoms are found in Antarctic waters (Figure 7). In Antarctica, diatom distribution is affected most profoundly by the seasonal progression of sea ice. This is especially true for large diatoms, because pore spaces within the ice are small and diatoms larger than approx. 30-40 microns are easily crushed in the freezing process. As such, the largest diatoms in Antarctica must survive beneath the sea ice – deep in the water column, in benthic and epibenthic habitats, or in the permanently open ocean beyond the sea ice. The 'giant' benthics also seem to be more cosmopolitan than the planktonic diatoms, with many occupying littoral regions in temperate and tropical latitudes as well as the icy coasts of Antarctica. *Arachnoidiscus japonicus* is more commonly found in the tropical Pacific and has only recently been found in Antarctica (Al-Handal & Wulff, 2008). Its size range in Antarctica appears to exceed descriptions elsewhere and may reflect an algal version of the 'gigantism' seen in other Antarctic marine organisms (Woods et al., 2009)!

The common Antarctic diatoms species *Corethron criophilum* and *C. pennatum* are exquisite examples for integrated mechanics on the micro- and nanoscale (Figures 3-5). Their structure and function is of high interest to nanotechnologists (Gebeshuber & Crawford, 2006; Gebeshuber, 2009). *C. criophilum* and *C. pennatum* exist as single cells (i.e., they do not form colonies) with two different valves per cell (heterovalvy). One hemispheric valve has a set of long spines that are attached to the valve at a series of sockets on the rim of the valve (Figure 8 right). The other valve has similar spines, but alternating with them is a series of finer hooked spines (Figure 8 left). The spines can move to a degree in the socket, but the position in which they are found in the mature, independent cell is not where they are formed. The process of new valve formation in

these diatoms is complicated (Crawford & Hinz, 1995; Crawford, Hinz & Honeywill, 1998). The spines are formed along with the new valves, within, and protected by the cylinder of the two sets of girdle bands (Figure 9). During the cell division cycle, the cell elongates greatly, forming very many girdle bands as it does so, thus creating a long space between the two sibling cells when they are complete. This space can accommodate the formation of the long spines. When the new cells are mature, they expand and pull the girdle cylinder away from the base of the spines and allow the spines to swing out to adopt their final position (Figure 10). In doing this they move past a click-stop that must be so structured as to ensure: a) that there are two positions in which the base of the spines can lie, b) that the first of these, lying beneath the girdle-band cylinder is unstable so that when released by the girdle cylinder it moves to the second position, c) that the final position of all three spine arrays is stable and d) that the orientation of all three arrays of spines differs (Richard Crawford, personal communication). This unique case of a click-stop mechanism in rigid micromechanical parts is a 'best practice' example for the connection of structure with function in nature.

The Antarctic diatom *Corethron* serves as inspiration for micro- and nanotechnology, where it is of paramount importance to come up with novel ideas on how to fabricate three-dimensional structures from two-dimensional structures, by unfolding and subsequently fixing them (Gebeshuber & Crawford, 2006; Gebeshuber et al., 2009). Imagine flat *Corethron*-inspired MEMS and NEMS that would unfold, expand and fix themselves in their predetermined final three-dimensional structure on demand!

One common characteristic of diatom shells is their fairly uniform pores (Figure 2). One potential use for these is to act as a selective filter, screening out larger molecules and letting smaller ones through (Figure 11).

Diatoms, due to their intricate structure at many size scales, have a huge surface area for their size. This makes them sensitive to tiny amounts of adsorbed gas molecules. Since their optical properties change when they adsorb gases, some experimental detectors of low levels of dangerous gases have been made from them (Lettieri et al., 2008). Various molecules can be bound to the surface of diatom shells, where they can react with other molecules and form a whole "lab on a diatom" (De Stefano et al., 2009).

One of the most advanced applications of diatoms is to turn their silica shells into other substances without changing their shapes. This is done by cooking them at high temperatures in a vapor containing atoms other than silicon. In this situation, when a silicon atom vaporizes from the shell, it is rapidly replaced by the alternative atom, so that the whole structure remains intact, not quite the Star Trek Replicator, but getting close (Drum & Gordon, 2003). The process is analogous to altering a brick building one brick at a time, replacing, say, each clay brick by a gold brick. The forces between adjacent atoms provide the "mortar". With this approach, ceramic (Dickerson et al., 2005) and metallic alloy (Sandhage & Bao, 2008) "diatoms" have been made. Silica is $\text{Si}(\text{OH})_4$, and a similar process removes the oxygen and hydrogen atoms, leaving pure silicon (Si). An all silicon "diatom" can possibly be used to create new three dimensional computers (Bao et al., 2007). Diatoms can also be used as molds to make nanostructures of other substances (Losic et al., 2007).

Genetic engineering of diatoms has already started with the complete sequencing of DNA in a few species (Genome Project, 2009b, c, a; Karthick, 2009) and direct manipulation of the genome (Kroth, 2007; Gordon et al., 2009).

The inherent bias towards benthic habitats means that Antarctica's giant diatoms are not hindered by heavy silicification and as such are far more robust than their tropical counterparts (eg. *Ethmodiscus* spp.). Whether or not large, heavily silicified shells have significant oil production by the cells is not yet known, though large Antarctic diatoms may play a role in global cooling through carbon sequestration (Pollock, 1997), to which they significantly contribute (DiTullio et al., 2000; Grigorov, Pearce & Kemp, 2002). Let's then return to the problem of a sustainable source of gasoline, which might be based on Antarctic diatoms.

The USA leads in gasoline consumption with an average of 10 barrels per person per year (StateMaster.com, 2009). Suppose we could genetically engineer giant Antarctic diatoms to secrete say 25% of their volume in oil per day, while living inside specially designed solar panels (Ramachandra et al., 2009). The largest Antarctic diatoms are nearly all epibenthic ones and the largest centric is *Arachnoidiscus* spp. (typically 0.4 to 1 mm). The volume of one giant 2 millimeter wide diameter subtropical and tropical species *Ethmodiscus rex* or *E. gazellae* cell is about 4 cubic millimeters (Villareal et al., 2007), so a big Antarctic *Arachnoidiscus* would about 1 cubic millimeter. Ten barrels converts to 1.64 billion cubic millimeters (1.64×10^9 mm³). This volume of oil could then possibly be produced by 20 million cells in the course of a year. A double layer of 1 million *Arachnoidiscus* cells of 1 mm diameter would occupy 10 square meters. While these cells have large vacuoles occupying perhaps 99% of their volume (Woods & Villareal, 2008), other diatoms do not and can have up to 85% of their volume as oils or lipids (Ramachandra et al., 2009). Furthermore, we might be able to shrink vacuole sizes by deliberate selection (Gordon, 1996). This then gives a rough calculation of how much area would need to be covered by gasoline secreting solar panels for each person. The production per cell might be less, but the number of layers of cells could be more, and there is a wide range of size of cells and approximately 100,000 species of diatoms (Fourtanier & Kocielek, 2009) to choose from, so if we could get diatom secretion of their oil to work, it might be practical. The advantages of solar panels are that they may be placed on roof tops, walls, deserts, and other generally unproductive areas, don't compete with farming for food, and may be widely distributed, eliminating transportation costs to deliver gasoline.

In summary, diatom nanotechnology may prove to be both a profitable and fundamental endeavor. Antarctic diatoms (WynnWilliams, 1996), including those in inland lakes (Jones, 1996; Laybourn-Parry & Pearce, 2007), come from a unique environment that is high in biodiversity (Brandt, 2005), and may be well worth our increased attention.

References

- Abdel-aal, H.A. & Ille C. Gebeshuber (2010). Special issue on: "Design in Nature". *International Journal of Design Engineering (IJDE)*, in preparation.
- Al-Handal, A.Y. & A. Wulff (2008). Marine epiphytic diatoms from the shallow sublittoral zone in Potter Cove, King George Island, Antarctica. *Botanica Marina* 51(5), 411-435.
- Allison, D.P., Y.F. Dufréne, M.J. Doktycz & M. Hildebrand (2008). Biomineralization at the nanoscale: learning from diatoms. *Methods in Cell Biology* 90, 61-86.
- Bao, Z., M.R. Weatherspoon, S. Shian, Y. Cai, P.D. Graham, S.M. Allan, G. Ahmad, M.B. Dickerson, B.C. Church, Z. Kang, H.W. Abernathy III, C.J. Summers, M. Liu & K.H. Sandhage (2007). Chemical reduction of three-dimensional silica micro-assemblies into microporous silicon replicas. *Nature* 446, 172-175.
- Bozarth, A., U.G. Maier & S. Zauner (2009). Diatoms in biotechnology: modern tools and applications. *Applied Microbiology and Biotechnology* 82(2), 195-201.
- Brandt, A. (2005). Evolution of Antarctic biodiversity in the context of the past: the importance of the Southern Ocean deep sea. *Antarctic Science* 17(4), 509-521.
- Crawford, R.M. & F. Hinz (1995). The spines of the centric diatom *Corethron criophilum*: light-microscopy of vegetative cell division. *European Journal of Phycology* 30(2), 95-105.
- Crawford, R.M., F. Hinz & C. Honeywill (1998). Three species of the diatom genus *Corethron* *Castracane*: structure, distribution and taxonomy. *Diatom Research* 13(1), 1-28.
- Cunningham, W.L. & A. Leventer (1998). Diatom assemblages in surface sediments of the Ross Sea: relationship to present oceanographic conditions. *Antarctic Science* 10(2), 134-146.
- Davis, J.S. (1972). Survival records in the algae, and the survival role of certain algae pigments, fat, and mucilaginous substances. *The Biologist* 54(2), 52-93.
- De Stefano, L., L. Rotiroti, M. De Stefano, A. Lamberti, S. Lettieri, A. Setaro & P. Maddalena (2009). Marine diatoms as optical biosensors. *Biosens. Bioelectron.* 24(6), 1580-1584.
- Dickerson, M.B., R.R. Naik, P.M. Sarosi, G. Agarwal, M.O. Stone & K.H. Sandhage (2005). Ceramic nanoparticle assemblies with tailored shapes and tailored chemistries via biosculpting and shape-preserving inorganic conversion. *Journal of Nanoscience and Nanotechnology* 5(1), 63-67.
- DíTullio, G.R., J.M. Grebmeier, K.R. Arrigo, M.P. Lizotte, D.H. Robinson, A. Leventer, J.B. Barry, M.L. VanWoert & R.B. Dunbar (2000). Rapid and early export of *Phaeocystis antarctica* blooms in the Ross Sea, Antarctica. *Nature* 404(6778), 595-598.

- Drum, R.W. & R. Gordon (2003). Star Trek replicators and diatom nanotechnology. *TibTech (Trends in Biotechnology)* 21(8), 325-328.
- Dupas, C., P. Houdy & M. Lahmani (2007). *Nanoscience: Nanotechnologies and Nanophysics*. Berlin, Springer-Verlag.
- Edlund, M.B. (2009). Diatoms in philately: postage stamps depicting diatoms. In: *20th North American Diatom Symposium, Iowa Lakeside Laboratory 23-27 September, 2009, and a special celebration of the life of Charles W. Reimer*. Eds.: M.B. Edlund & S. Spaulding. <http://www.northamericadiatomsymposium.org/pdfs/Edlund.pdf>.
- Fourtanier, E. & J.P. Kociolek. (2009). Catalogue of Diatom Names. <http://research.calacademy.org/research/diatoms/names/>
- Fryxell, G.A. (1994). Planktonic marine diatom winter stages: Antarctic alternatives to resting spores. *Memoirs of the San Francisco Academy of Sciences* 17, 437-448.
- Gazit, E. (2007). *Plenty of Room for Biology at the Bottom: An Introduction to Bionanotechnology*. Singapore, World Scientific.
- Gebeshuber, I.C. & R.M. Crawford (2006). Micromechanics in biogenic hydrated silica: hinges and interlocking devices in diatoms. *Proceedings of the Institution of Mechanical Engineers Part J-Journal of Engineering Tribology* 220(J8), 787-796.
- Gebeshuber, I.C. (2009). Nanoelectromechanical systems (NEMS) – Introduction, application and challenges of nanoelectromechanical systems. <http://www.azonano.com/details.asp?ArticleId=2465>
- Gebeshuber, I.C., H. Stachelberger, B.A. Ganji, D.C. Fu, J. Yunas & B.Y. Majlis (2009). Exploring the innovational potential of biomimetics for novel 3D MEMS. *Journal of Advanced Materials Research* 74, 265-268.
- Genome Project. (2009a). *Cyclotella meneghiniana*. <http://www.ncbi.nlm.nih.gov/sites/entrez?Db=genomeprj&cmd=ShowDetailView&TermToSearch=32153>
- Genome Project. (2009b). *Thalassiosira pseudonana*. <http://www.ncbi.nlm.nih.gov/sites/entrez?Db=genomeprj&cmd=ShowDetailView&TermToSearch=9562>
- Genome Project. (2009c). *Phaeodactylum tricornutum*. <http://www.ncbi.nlm.nih.gov/sites/entrez?Db=genomeprj&cmd=ShowDetailView&TermToSearch=13153>
- Goodsell, D.S. (2004). *Bionanotechnology, Lessons from Nature*. Hoboken, NJ, Wiley-Liss.
- Gordon, R. (1987). A retaliatory role for algal projectiles, with implications for the mechanochemistry of diatom gliding motility. *J. Theor. Biol.* 126, 419-436.
- Gordon, R. & B.D. Aguda (1988). Diatom morphogenesis: natural fractal fabrication of a complex microstructure. In: *Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, Part 14: Cardiology and Imaging, 4-7 Nov. 1988, New Orleans, LA, USA*. Eds.: G. Harris & C. Walker. New York, Institute of Electrical and Electronics Engineers: 273-274.
- Gordon, R. (1996). Computer controlled evolution of diatoms: design for a compustat. *Nova Hedwigia* 112(Festschrift for Prof. T.V. Desikachary), 213-216.
- Gordon, R., F.A.S. Sterrenburg & K. Sandhage (2005). A Special Issue on Diatom Nanotechnology. *Journal of Nanoscience and Nanotechnology* 5(1), 1-4.
- Gordon, R. (2008). The Glass Menagerie: Diatom Nanotechnology and Its Implications for Multi-Scale Manufacturing and Oil Production. In: *2nd International Conference on Multi-Scale Structures and Dynamics of Complex Systems: Processes & Forces for Creation of Designer Materials with Multi-Scale Structures, 4-5 September, 2008, Bangalore, India*. Eds.: J.S. Raut & R. Venkataraghavan. Bangalore, Unilever.
- Gordon, R., D. Losic, M.A. Tiffany, S.S. Nagy & F.A.S. Sterrenburg (2009). The Glass Menagerie: diatoms for novel applications in nanotechnology [invited]. *Trends in Biotechnology* 27(2), 116-127.
- Gordon, R. (2010). Diatoms and nanotechnology: early history and imagined future as seen through patents [Invited]. In: *The Diatoms: Applications for the Environmental and Earth Sciences*. Eds.: J.P. Smol & E.F. Stoermer. Cambridge, Cambridge University Press. 2nd: in press.
- Grigorov, I., R.B. Pearce & A.E.S. Kemp (2002). Southern Ocean laminated diatom ooze: mat deposits and potential for palaeo-flux studies, ODP leg 177, Site 1093. *Deep-Sea Research Part II-Topical Studies in Oceanography* 49(16), 3391-3407.
- Helbling, E.W., B.E. Chalker, W.C. Dunlap, O. Holmhansen & V.E. Villafane (1996). Photoacclimation of Antarctic marine diatoms to solar ultraviolet radiation. *Journal of Experimental Marine Biology and Ecology* 204(1-2), 85-101.
- Hernando, M.P. & G.A. Ferreyra (2005). The effects of UV radiation on photosynthesis in an Antarctic diatom (*Thalassiosira* sp.): Does vertical mixing matter? *Journal of Experimental Marine Biology and Ecology* 325(1), 35-45.
- Hollibaugh, J.T., D.L.R. Seibert & W.H. Thomas (1981). Observations on the survival and germination of resting spores of three *Chaetoceros* (Bacillariophyceae) species. *Journal of Phycolgy* 17, 1-9.
- Janech, M.G., A. Krell, T. Mock, J.S. Kang & J.A. Raymond (2006). Ice-binding proteins from sea ice diatoms (Bacillariophyceae). *Journal of Phycolgy* 42(2), 410-416.
- Jones, V.J. (1996). The diversity, distribution and ecology of diatoms from Antarctic inland waters. *Biodiversity and Conservation* 5(11), 1433-1449.
- Karthick, B. (2009). Genome sequencing of cells that live inside glass cages reveals their past history. *Current Science* 96(3), 334-337.
- Kemp, K.D. (2009). MicroLife Services, Specialist in diatom microslides and diatom arrangements. <http://www.diatoms.co.uk>

- Krell, A., B. Beszteri, G. Dieckmann, G. Glockner, K. Valentin & T. Mock (2008). A new class of ice-binding proteins discovered in a salt-stress-induced cDNA library of the psychrophilic diatom *Fragilariopsis cylindrus* (Bacillariophyceae). *European Journal of Phycology* 43(4), 423-433.
- Kröger, N. & N. Poulsen (2008). Diatoms: from cell wall biogenesis to nanotechnology. *Annu. Rev. Genet.* 42, 83-107.
- Kroth, P., R. León, A. Gaván & E. Fernández (2007). Molecular biology and the biotechnological potential of diatoms. *Advances in Experimental Medicine & Biology* 616, 23-33.
- Kroth, P.G. (2007). Genetic transformation: a tool to study protein targeting in diatoms. *Methods Mol Biol* 390, 257-268.
- Laughlin, R.B. (2005). *A Different Universe: Reinventing Physics from the Bottom Down*. New York, Basic Books.
- Laybourn-Parry, J. & D.A. Pearce (2007). The biodiversity and ecology of Antarctic lakes: models for evolution. *Philosophical Transactions of the Royal Society B-Biological Sciences* 362(1488), 2273-2289.
- Lettieri, S., A. Setaro, L. De Stefano, M. De Stefano & P. Maddalena (2008). The gas-detection properties of light-emitting diatoms. *Advanced Functional Materials* 18(8), 1257-1264.
- Ligowski, R., M. Godlewski & A. Łukowski (1992). Sea ice diatoms and ice edge planktonic diatoms at the northern limit of the Weddell Sea pack ice. *Proc. NIPR Symp. Polar Biol.* 5, 9-20.
- Link, R.J. & D. Zimmerman (2007). An approach for model updating of a multiphysics MEMS micromirror. *Journal of Dynamic Systems Measurement and Control-Transactions of the ASME* 129(3), 357-366.
- Losic, D., J.G. Mitchell, R. Lal & N.H. Voelcker (2007). Rapid fabrication of micro- and nanoscale patterns by replica molding from diatom biosilica. *Advanced Functional Materials* 17(14), 2439-2446.
- Matthias Burba. (2008). Diatoms (from an 1891 J.D. Möller arrangement) (65X). <http://www.nikonsmallworld.com/moreinfo.php?grouping=year&year=2008&imagepos=50>
- Medlin, L.K. & J. Priddle (1990). *Polar Marine Diatoms*. Cambridge, British Antarctic Survey, Natural Environment Research Council.
- Nagy, S.S. (2002). Montana Diatoms. <http://montanadiatoms.tripod.com/>
- Papazoglou, E.S. & A. Parthasarathy (2007). *BioNanotechnology*, Morgan and Claypool Publishers.
- Parkinson, J. & R. Gordon (1999). Beyond micromachining: the potential of diatoms. *Trends Biotechnol (Tibtech)* 17(5), 190-196.
- Pollock, D.E. (1997). The role of diatoms, dissolved silicate and Antarctic glaciation in glacial/interglacial climatic change: a hypothesis. *Global and Planetary Change* 14(3-4), 113-125.
- Ramachandra, T.V., D.M. Mahapatra, Karthick B. & R. Gordon (2009). Milking diatoms for sustainable energy: biochemical engineering versus gasoline-secreting diatom solar panels [invited]. *Industrial & Engineering Chemistry Research* 48(19, Complex Materials II special issue, October), 8769-8788.
- Rehm, B.H.A. (2006). *Microbial Bionanotechnology: Biological Self-Assembly Systems and Biopolymer-Based Nanostructures*, Taylor & Francis.
- Reisner, D.E., Ed. (2008). *Bionanotechnology: Global Prospects*. Boca Raton, CRC.
- Renugopalakrishnan, V. & R.V. Lewis, Eds. (2006). *Bionanotechnology: Proteins to Nanodevices*, Springer.
- Ruddiman, W.E. (2005). How did humans first alter global climate? *Scientific American* 292(3), 46-53.
- Sandhage, K.H. & Z. Bao (2008). *Methods of fabricating nanoscale-to-microscale structures [United States Patent Application US 2008/0038170 A1]*.
- Sargent, E.H. (2005). *The Dance of Molecules: How Nanotechnology is Changing Our Lives*. Toronto, Viking Canada.
- Scharek, R. (2010). Antarctic diatoms, a biological view. In: *Antarctica. Time of Change*. Barcelona, ACTAR / Arts Santa Mònica.
- Skerratt, J.H., A.D. Davidson, P.D. Nichols & T.A. McMeekin (1998). Effect of UV-B on lipid content of three antarctic marine phytoplankton. *Phytochemistry* 49(4), 999-1007.
- StateMaster.com. (2009). Energy Statistics > Gasoline Consumption (per capita) (most recent) by state. http://www.statemaster.com/graph/ene_gas_con_percap-energy-gasoline-consumption-per-capita
- Sterrenburg, F.A.S., R. Gordon, M.A. Tiffany & S.S. Nagy (2007). Diatoms: living in a construal environment. In: *Algae and Cyanobacteria in Extreme Environments. Series: Cellular Origin, Life in Extreme Habitats and Astrobiology, Vol. 11*. Eds.: J. Seckback. Dordrecht, The Netherlands, Springer: 141-172.
- Villareal, T.A., R.M.L. McKay, M.M.D. Al-Rshaidat, R. Boyanapalli & R.M. Sherrell (2007). Compositional and fluorescence characteristics of the giant diatom *Ethmodiscus* along a 3000 km transect (28 degrees N) in the central North Pacific gyre. *Deep-Sea Research Part I-Oceanographic Research Papers* 54(8), 1273-1288.
- Weatherspoon, M.R., M.B. Dickerson, G. Wang, Y. Cai, S. Shian, S.C. Jones, S.R. Marder & K.H. Sandhage (2007). Thin, conformal, and continuous SnO₂ coatings on three-dimensional biosilica templates through hydroxy-group amplification and layer-by-layer alkoxide deposition. *Angewandte Chemie-International Edition* 46(30), 5724-5727.
- Witkowski, A., C. Riaux-Gobin & G. Daniszewska-Kowalczyk (2010). Diatom (Bacillariophyceae) flora of the Austral Islands Kerguelen, with consideration on biogeographic distribution of established and newly described taxa. *Iconogr. Diatomol.*, submitted.
- Woods, H.A., A.L. Moran, C.P. Arango, L. Mullen & C. Shields (2009). Oxygen hypothesis of polar gigantism not supported by performance of Antarctic pycnogonids in hypoxia. *Proc. R. Soc. B-Biol. Sci.* 276(1659), 1069-1075.
- Woods, S. & T.A. Villareal (2008). Intracellular ion concentrations and cell sap density in positively buoyant oceanic phytoplankton. *Nova Hedwigia(Suppl.)* 133, 131-145.

Wulff, A., M.Y. Roleda, K. Zacher & C. Wiencke (2008). Exposure to sudden light burst after prolonged darkness: a case study on benthic diatoms in Antarctica. *Diatom Research* 23(2), 519-532.

WynnWilliams, D.D. (1996). Antarctic microbial diversity: The basis of polar ecosystem processes. *Biodiversity and Conservation* 5(11), 1271-1293.

Zgurovskaya, L.N. (1977). Effect of addition of nutrients on growth of spores and division of planktonic algae from bottom sediments. *Okeanologiya* 17(1), 119-122.

Zielinski, U. & R. Gersonde (1997). Diatom distribution in Southern Ocean surface sediments (Atlantic sector): Implications for paleoenvironmental reconstructions. *Palaeogeography Palaeoclimatology Palaeoecology* 129(3-4), 213-250.

Richard Gordon, a theoretical biologist, got hooked on diatoms because of the fascinating movement of the colonial diatom *Bacillaria paradoxa*. He has tried to understand the motility mechanism and the morphogenesis of the diatom shell, meanwhile inadvertently starting the field of diatom nanotechnology. His other research includes morphogenesis of early vertebrate embryos, detection of early breast cancer, and halting of the HIV/AIDS epidemic.

Andrzej Witkowski is a geologist and oceanographer by training who got involved in diatom studies while working on his Ph.D. on fossilization processes of siliciclastic microbial mats. Since then his research has focused on paleo aspects of marine diatoms and species composition, biodiversity and biogeography of marine littoral diatoms worldwide. Recently he has investigated marine littoral diatoms of the South African and Kerguelen Islands and some parts of Antarctica. He is Editor of Diatom Monographs.

Ile C. Gebeshuber is an experimental physicist who has been working with diatoms since 1999 when she was the first to image live diatoms *in situ* under water with nanometer resolution and found self-repair properties of diatom adhesives, which make them interesting for man-made high-tech underwater glues. In 2004 she founded the field of diatom tribology, where diatoms are used as inspiration for the development of man-made micro- and nanoelectromechanical machines (MEMS and NEMS). The amazing click-stop mechanism in the Antarctic diatoms *Corethron pennatum* and *C. criophilum* inspires us to build the tiniest optimized MEMS structures that unfold and are then irreversibly fixed.

Claire Allen is a palaeoceanographer at the British Antarctic Survey, currently working within the 'Quaternary Sediments' work package of the 'Chemistry and Past Climate' research programme. She uses diatoms preserved in marine sediments to reconstruct past changes in the ocean and climate of Antarctica to better understand Antarctica's role in the global climate system. Claire has participated in six Antarctic science cruises throughout the Antarctic Peninsula and Scotia Sea, collecting water and sediment samples to help constrain the ecological significance of Southern Ocean diatom records.

VICENÇ ALTAIÓ
JOSEP PERELLÓ
ALICIA CHILLIDA
MERCEDES MASÓ
PATRICIA HOMS
MIREYA MASÓ
MARA DIERSSEN
VÍCTOR GÓMEZ PIN
ENRIC PUIG PUNYET
TED SCAMBOS
JOSÉ MANUEL FORTUÑO
RENAME SCHAREK
RICHARD GORDON
ANDRZEJ WITKOWSKI
ILLE CHRISTINE GEBESHUBER
CLAIRE S. ALLEN
MARTA MALÉ
LUIS FRAGUADA



Generalitat de Catalunya-
Departament de Cultura
i Mitjans de Comunicació

